
Identification of protease exosite-interacting peptides that enhance substrate cleavage kinetics.

Journal: Biol Chem

Publication Year: 2012

Authors: Abeer M Jabaiah, Jennifer A Getz, Witold A Witkowski, Jeanne A Hardy, Patrick S Daugherty

PubMed link: 22944693

Public Summary:

Scientific Abstract:

Many peptidases are thought to require non-active site interaction surfaces, or exosites, to recognize and cleave physiological substrates with high specificity and catalytic efficiency. However, the existence and function of protease exosites remain obscure owing to a lack of effective methods to identify and characterize exosite-interacting substrates. To address this need, we modified the cellular libraries of peptide substrates (CLiPS) methodology to enable the discovery of exosite-interacting peptide ligands. Invariant cleavage motifs recognized by the active sites of thrombin and caspase-7 were displayed on the outer surface of bacteria adjacent to a candidate exosite-interacting peptide. Exosite peptide libraries were then screened for ligands that accelerate cleavage of the active site recognition motif using two-color flow cytometry. Exosite CLiPS (eCLiPS) identified exosite-binding peptides for thrombin that were highly similar to a critical exosite interaction motif in the thrombin substrate, protease-activated receptor 1. Protease activity probes incorporating exosite-binding peptides were cleaved ten-fold faster than substrates without exosite ligands, increasing their sensitivity to thrombin activity in vitro. For comparison, screening with caspase-7 yielded peptides that modestly enhanced (two-fold) substrate cleavage rates. The eCLiPS method provides a new tool to profile the ligand specificity of protease exosites and to develop improved substrates.

PNAS Lens Free Article Link:



Source URL: <https://www.cirm.ca.gov/about-cirm/publications/identification-protease-exosite-interacting-peptides-enhance-substrate>